

Features of the reduction of graphene from graphene oxide

V. A. Labunov ¹,

L. V. Tabulina ²,

I. V. Komissarov ³,

D. V. Grapov ⁴,

E. L. Prudnikova ⁵,

Yu. P. Shaman (Foreign) ⁶,

S. A. Basaev (Foreign) ⁷,

A. A. Pavlov (Foreign) ⁸

2017

1, 2, 3, 4, 5 Belarusian State University of Informatics and Radioelectronics, P. Browka 6, 220013 Minsk, Belarus

6, 7 Foreign (NPK Technological Center, National Research University of Electronic Technology (MIET), Zelenograd, Moscow, Russia)

8 Foreign (Institute of Microelectronic Nanotechnologies, Russian Academy of Sciences, Moscow, Russia)

Keywords: Graphene, graphene oxide, reduction of graphene oxide, Hummers method nanostructures.

Abstract: Features of the reduction of graphene from graphene oxide in media containing hydrazine hydrate, ethylene glycol, and hydrogen are studied. X-ray energy dispersive spectroscopy, Raman spectroscopy, and

scanning electron microscopy data indicate that this process proceeds through the high-temperature annealing of graphene oxide in a hydrogen environment.

Published in:

Russian Journal of Physical Chemistry A – June 2017, Volume 91, Issue 6, pp. 1088–1092. – <https://doi.org/10.1134/S0036024417060140>.

Read More:

<https://link.springer.com/article/10.1134/S0036024417060140>.

© Pleiades Publishing, Ltd. 2017.